

SYMMETRIC ALCOVED POLYTOPES

ANNETTE WERNER AND JOSEPHINE YU

ABSTRACT. Generalized alcoved polytopes are polytopes whose facet normals are roots in a given root system. We call a set of points in an alcoved polytope a generating set if there exists no strictly smaller alcoved polytope containing it. The type A alcoved polytopes are precisely the convex polytopes that are also tropically convex. In this case the tropical generators form a generating set. We show that for any root system other than F_4 , every alcoved polytope invariant under the natural Weyl group action has a generating set of cardinality equal to the Coxeter number of the root system.

1. INTRODUCTION

In this paper we investigate alcoved polytopes which are symmetric under the action of the Weyl group from the viewpoint of tropical convexity. Replacing addition by tropical addition and multiplication by tropical multiplication, one can define tropically convex subsets of finite dimensional real vector spaces. Develin and Sturmfels showed that tropical polytopes have a natural polyhedral cell decomposition such that the cells are tropically convex and classically convex at the same time [De-St]. Such sets are named *polytropes* by Joswig and Kulas [Jo-Ku]. Every polytrope can be written as an intersection of halfspaces given by root hyperplanes for a root system of type A [De-St].

This paper grew out of an attempt to generalize notions of tropical convexity to other types of root systems. The polytropes, which are building blocks of tropically convex sets, have a characterization that natural generalizes to other root systems: they are alcoved polytopes of type A studied by Lam and Postnikov [La-Po]. We consider them in a slightly more general situation than in [La-Po], since we also allow non-integral coefficients. Any d -dimensional alcoved polytope of type A is the tropical convex hull of $d + 1$ vertices [Jo-Ku]. We are interested in generating sets for alcoved polytopes associated to arbitrary root systems. Here a set of points in an alcoved polytope P is called a generating set if P is the smallest alcoved polytope containing it.

Our main result states that every alcoved polytope for an irreducible and reduced root system Φ not of type F_4 that is symmetric under the action of the Weyl group can be generated by h vertices, where h is the Coxeter number of Φ . We also discuss the case F_4 , where a symmetric alcoved polytope may need up to 24 generators while the Coxeter number is 12.

In the A_n -case however, *every* alcoved polytope has a generating set of size $n + 1$ which is the Coxeter number, regardless of symmetry. We give a short proof of this fact. The symmetry assumption cannot be dropped in general, since there exists a non-symmetric

alcoved polytope of type D_4 requiring eight generators, whereas the Coxeter number of D_4 is six.

2. TROPICAL CONVEXITY AND ALCOVED POLYTOPES

We begin by recalling some facts on tropical convexity. Let $(\mathbb{R}, \oplus, \odot)$ be the min-plus tropical semiring. The space \mathbb{R}^n together with componentwise addition \oplus is a semimodule under the semiring $(\mathbb{R}, \oplus, \odot)$, if we put $a \odot (x_1, \dots, x_n) = (a + x_1, \dots, a + x_n)$ for $a \in \mathbb{R}$ and $(x_1, \dots, x_n) \in \mathbb{R}^n$.

Definition 2.1 ([De-St]). A subset S of \mathbb{R}^n is called *tropically convex* if for all x, y in S and for all $\mu, \nu \in \mathbb{R}$ the element $(\mu \odot x) \oplus (\nu \odot y)$ is also contained in S . The *tropical convex hull* of a subset V in \mathbb{R}^n is the smallest tropically convex set containing V .

Note that any tropically convex subset of \mathbb{R}^n containing x also contains $x + (r, \dots, r) = r \odot x$ for all $r \in \mathbb{R}$. Therefore we project tropically convex subsets of \mathbb{R}^n to $\mathbb{R}^n/\mathbb{R}(1, \dots, 1)$ and call their image in $\mathbb{R}^n/\mathbb{R}(1, \dots, 1)$ also tropically convex.

In [De-St, Theorem 15] it is shown that the tropical convex hull of a finite subset of $\mathbb{R}^n/\mathbb{R}(1, \dots, 1)$ admits a polyhedral cell decomposition. These bounded cells are at the same time tropically convex and convex in the ordinary sense. Such a set is also called a *polytrope*; see [Jo-Ku].

Tropical convexity is closely related to root systems of type A in the following way. Let U be the subspace $\{(x_1, \dots, x_n) : \sum_i x_i = 0\}$ of \mathbb{R}^n . The quotient map

$$U \rightarrow \mathbb{R}^n/\mathbb{R}(1, \dots, 1)$$

is an isomorphism of real vector spaces. Hence the restriction of the canonical scalar product on \mathbb{R}^n to U induces a scalar product $\langle \cdot, \cdot \rangle$ on $\mathbb{R}^n/\mathbb{R}(1, \dots, 1)$. Consider the elements $a_{ij} = e_i - e_j$ in $\mathbb{R}^n/\mathbb{R}(1, \dots, 1)$, where e_i denotes the canonical unit vectors. Then $\Phi = \{a_{ij} : i \neq j\}$ is a root system of type A_{n-1} in $\mathbb{R}^n/\mathbb{R}(1, \dots, 1)$.

Lemma 2.2. *For any polytrope P in $\mathbb{R}^n/\mathbb{R}(1, \dots, 1)$ there are non-negative real constants c_{ij} such that*

$$P = \bigcap_{i \neq j} \{x \in \mathbb{R}^n/\mathbb{R}(1, \dots, 1) : \langle a_{ij}, x \rangle \leq c_{ij}\}.$$

Proof. By [De-St, Lemma 10 and Proposition 18], polytropes are precisely the bounded intersections of half-spaces of the form $\{x = (x_1, \dots, x_n) \in \mathbb{R}^n/\mathbb{R}(1, \dots, 1) : x_i - x_j \leq c_{ij}\}$ for $i \neq j$ and $c_{ij} \in \mathbb{R}$, which implies our claim. \square

Hence polytropes look like alcoved polytopes of type A , as defined in [La-Po]. To be precise, let V be a real vector space endowed with a scalar product $\langle \cdot, \cdot \rangle$, and let Φ be any irreducible and reduced root system in V . By t we denote the type of the corresponding Dynkin diagram, i.e. t is equal to A_n for $n \geq 1$, B_n , C_n for $n \geq 2$, D_n for $n \geq 4$ or to E_6 , E_7 , E_8 , F_4 , or G_2 . Moreover, let W be the Weyl group of V , i.e. W is the finite group of orthogonal endomorphisms of V generated by the reflections at the hyperplanes $H_a = \{x \in V : \langle a, x \rangle = 0\}$ perpendicular to the roots.

Definition 2.3. A *generalized alcoved polytope of type t* is a bounded intersection of root halfspaces of the form

$$\bigcap_{a \in \Psi} \{x \in V : \langle a, x \rangle \leq c_a\},$$

where Ψ is a subset of a root system Φ of type t and $c_a \in \mathbb{R}$.

Thus a generalized alcoved polytope is a bounded subset of V defined as an intersection of root halfspaces. Alternatively, we can describe it as a polytope whose set of facet normal vectors is contained in Φ . If the right hand side values c_a are all integers, a generalized alcoved polytope we defined is an alcoved polytope from [La-Po]. Payne [Pa] showed that alcoved polytopes have a Koszul property. For simplicity we will refer to “generalized alcoved polytopes” simply as “alcoved polytopes”.

Some examples of alcoved polytopes include associahedra (see [Jo-Ku] and references therein), cubes, cross polytopes, and the 24-cell as a degenerate F_4 alcoved polytope.

3. GENERATING SETS OF ALCOVED POLYTOPES

We will define generating sets of alcoved polytopes and prove results about their cardinality for type A . Besides, we discuss an interesting example for type D_4 .

Definition 3.1. Let Φ be a root system of type t in V , and let P be an alcoved polytope of type t in V . A subset $\Sigma \subset V$ is called a *generating set* of P if P is the smallest alcoved polytope in V containing Σ .

Let P be an alcoved polytope for the root system Φ of type A_{n-1} as in Lemma 2.2. Since alcoved polytopes are tropically convex, every subset of P whose tropical convex hull is P is a generating set. However not every set of generators of P in the sense of Definition 3.1 is a set of tropical generators. The tropical convex hull of a generating set may be strictly smaller than P . For example, consider the square with vertices $(0, 0, 0), (0, 0, 1), (0, 1, 0), (0, 1, 1)$ in $\mathbb{R}^3/\mathbb{R}(1, 1, 1)$. The two vertices $(0, 0, 1)$ and $(0, 1, 0)$ generate the square as an alcoved polytope, but their tropical convex hull is strictly smaller. In fact, the square cannot be generated tropically by two points, as the tropical convex hull of any two points is at most one dimensional in $\mathbb{R}^n/\mathbb{R}(1, \dots, 1)$.

For any bounded $S \subseteq V$ and $a \in \Phi$ the smallest halfspace of the form $\{x \in V : \langle a, x \rangle \leq c\}$ that contains S is $\{x \in V : \langle a, x \rangle \leq \sup_{s \in S} \langle a, s \rangle\}$, so the alcoved polytope generated by S is

$$\bigcap_{a \in \Phi} \{x \in V : \langle a, x \rangle \leq \sup_{s \in S} \langle a, s \rangle\}.$$

On the other hand, for an alcoved polytope P , a subset $S \subset P$ is a generating set if and only if for every $a \in \Phi$ we have

$$\sup_{x \in P} \langle a, x \rangle = \sup_{x \in S} \langle a, x \rangle.$$

This can be rephrased as follows. Recall that a *support hyperplane* H of a polytope P is a hyperplane containing P on one side and having non-empty intersection with P . The intersection $H \cap P$ is the *face* of P supported by H .

Lemma 3.2. *A set $S \subseteq P$ is a generating set for P if and only if for any $a \in \Phi$ the support hyperplane of P perpendicular to a contains at least one point from S .*

It follows that there is a minimal cardinality generating set consisting of vertices of P only. If a point in a generating set lies in the relative interior of a non-vertex face, then the point can be replaced by a vertex of that face, as doing so only enlarges the set of support hyperplanes containing the point.

Question 3.3. *Our problem can be phrased as follows.*

- (1) *For each root system Φ , find a number k such that every alcoved polytope of type Φ has a generating set of size at most k .*
- (2) *More generally, let P be a polytope and \mathcal{H} be a set of support hyperplanes of P such that all facets of P are contained in \mathcal{H} . The containment may be strict. Find the minimal cardinality of a set $S \subset P$ such that every support hyperplane in \mathcal{H} contains at least one point from S .*

There is a nice solution for type A alcoved polytopes, as we will see in Proposition 3.6 below. The general problem seems computationally difficult, as it amounts to solving a set covering problem, which is known to be NP-complete in general.

Remark 3.4. For every root system Φ there exist alcoved polytopes which cannot be generated with less than $|\Phi|/\dim(V)$ vertices. For a fixed set of facet normals, the condition on the coefficients for the perpendicular hyperplanes to meet at a point is described by vanishing of certain determinants. Generic choices of coefficients would avoid this determinantal locus and give polytopes that are simple, i.e. no more than $\dim(V)$ facets can meet at one point. Therefore in this case we need at least $|\Phi|/\dim(V)$ vertices to support all facets.

Proposition 3.5. *Let t be the type of a root system Φ in a vector space of dimension 2. Then every alcoved polytope of type t can be generated by $|\Phi|/2$ many vertices.*

Proof. Let us cyclically order the roots using their positions in the plane. For any two support hyperplanes corresponding to two consecutive roots, there is a vertex supporting them, so $|\Phi|/2$ vertices are sufficient to support all root hyperplanes. \square

In [Jo-Ku, Theorem 12], it is shown that every alcoved polytope of type A_n is a tropical simplex, i.e. it has $d + 1$ tropical generators, where d is the dimension of the alcoved polytope (which might be smaller than n if the polytope lies in a root hyperplane). In particular it can be generated by $n + 1$ elements in the sense of Definition 3.1. The next proposition provides an easy proof of this fact. This does not imply the Joswig–Kulas result because the set of tropical generators may be strictly larger.

Proposition 3.6. *Every alcoved polytope of type A_n is generated by $n + 1$ elements.*

Proof. Let P be a nonempty alcoved polytope of type A_n defined by $x_i - x_j \leq c_{ij}$ for some $c_{ij} \in \mathbb{R}$ where $0 \leq i, j \leq n$ and $i \neq j$. By Lemma 3.2, we need to consider the support hyperplanes of P normal to the roots, so we may assume that all inequalities are tight, i.e. for each pair (i, j) , there is a point $p \in P$ such that $p_i - p_j = c_{ij}$. Then we

must have $c_{ik} - c_{jk} \leq c_{ij}$ for any distinct i, j, k . Otherwise, for any $p \in P$, $p_i - p_k = (p_i - p_j) + (p_j - p_k) \leq c_{ij} + c_{jk} < c_{ik}$, contradicting our assumption that the inequality $x_i - x_k \leq c_{ik}$ is tight for P .

For each $0 \leq k \leq n$, let $p^{(k)} = (c_{0k}, \dots, c_{k-1,k}, 0, c_{k+1,k}, \dots, c_{nk}) \in \mathbb{R}^{n+1}/\mathbb{R}(1, \dots, 1)$. For any $l \neq k$, the point $p^{(k)}$ lies on the hyperplanes $x_\ell - x_k = c_{\ell k}$. Since P is non empty, let $p \in P$ and we get $-c_{\ell k} \leq p_k - p_\ell \leq c_{k\ell}$. Hence $p_k^{(k)} - p_\ell^{(k)} = -c_{\ell k} \leq c_{k\ell}$. Moreover, for any pair $i \neq j$ distinct from k , we have $p_i^{(k)} - p_j^{(k)} = c_{ik} - c_{jk} \leq c_{ij}$ using the inequality seen above. Therefore the point $p^{(k)}$ lies in the polytope P . The $n+1$ points $p^{(0)}, p^{(1)}, \dots, p^{(n)}$ all lie in P and support all the root hyperplanes, so they form a generating set of size $n+1$. \square

Note that $n+1$ is the Coxeter number of the root system of type A_n . We will see below in Theorem 4.3 that almost all Weyl group symmetric alcoved polytopes can be generated by a set of vertices whose cardinality is the Coxeter number of the root system.

However, for types of root systems other than A_n this is no longer true if we drop the symmetry hypothesis, as seen in the following example.

Example 3.7. Consider the D_4 alcoved polytope in \mathbb{R}^4 defined by the inequalities given in the columns of the following matrix in homogeneous coordinates. For example, the first column represents the inequality $100 + x_1 + x_2 \geq 0$.

$$\begin{pmatrix} 100 & 97 & 96 & 95 & 96 & 98 & 95 & 98 & 96 & 98 & 96 & 96 & 98 & 98 & 99 & 99 & 95 & 96 & 95 & 96 & 95 & 99 & 95 & 100 \\ 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 & 0 & 0 & 0 & 0 & 1 & -1 & 1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & -1 & -1 & 1 & 0 & 0 & 0 & 0 & 1 & -1 & 1 & -1 & 0 & 0 & 0 & 0 & 1 & -1 & 1 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -1 & -1 & 1 & 1 & -1 & -1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & -1 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & -1 & -1 & 1 & 1 & -1 & -1 & 1 & -1 & -1 & 1 \end{pmatrix}$$

A computer check shows that this alcoved polytope requires eight generators, while the Coxeter number of D_4 is six. The polytope is simple and has f -vector $(96, 192, 120, 24, 1)$.

4. SYMMETRIC ALCOVED POLYTOPES

In this section we define symmetric alcoved polytopes and prove our main result stating that in all cases except F_4 they can be generated by a set of cardinality equal to the Coxeter number. Afterwards we discuss the case of F_4 -alcoved polytopes.

Let Φ be a root system contained in the Euclidean vector space (V, \langle, \rangle) and let W be the associated Weyl group.

Definition 4.1. We call an alcoved polytope in V of type t *symmetric* if it is invariant under the action of the Weyl group W .

Note that W acts on the faces of a symmetric alcoved polytope. Since every w in W maps the support hyperplane with respect to the root a to the support hyperplane with respect to the root $w(a)$, we can write a symmetric alcoved polytope as

$$P = \bigcap_{a \in \Phi} \{x \in V : \langle a, x \rangle \leq c_a\},$$

with $c_a = c_{w(a)}$ for all w in W and for all $a \in \Phi$.

Note that an alcoved polytope for a non-reduced root system Φ is also an alcoved polytope of the associated reduced root system Φ^{red} . Moreover, alcoved polytopes for

reducible root systems are products of alcoved polytopes for the irreducible components. Therefore we will from now on assume that Φ is irreducible and reduced.

Let us recall some basic facts about root systems from [Bou, Chapter VI]. For every root a we denote by s_a the corresponding reflection in V . Then for all roots b we have

$$s_a(b) = a - n(a, b)b, \quad \text{where } n(a, b) = 2 \frac{\langle a, b \rangle}{\langle b, b \rangle}$$

is a number between -3 and 3 in the reduced case.

Fix a chamber with respect to the hyperplane arrangement in V given by the hyperplanes $H_a = \{x \in V : \langle a, x \rangle = 0\}$. Let b_1, \dots, b_n be the roots perpendicular to the facets of the fixed chamber which lie on the same side as the chamber. They form a basis of Φ [Bou, VI, 1.5, Theorem 2]. Then the element

$$\omega = s_{b_1} \circ \dots \circ s_{b_n}$$

of the Weyl group is called a *Coxeter element*. It depends up to conjugation on the choice of the basis and on the choice of its ordering [Bou, V, 6.1, Proposition 1]. The order h of ω in the Weyl group is called the *Coxeter number* of Φ . By [Bou, V, 6.2, Theorem 1], we have

$$nh = |\Phi|$$

where $|\Phi|$ is the number of roots.

After choosing a basis b_1, \dots, b_n we also write $s_i = s_{b_i}$. Let Γ be the cyclic subgroup of W generated by $\omega = s_1 \circ \dots \circ s_n$. For all $i = 1, \dots, n$ let

$$\theta_i = s_n \circ s_{n-1} \circ \dots \circ s_{i+1}(b_i).$$

Proposition 4.2. [Bou, VI, 1.11, Proposition 33] *With the notation above, the root system Φ is a disjoint union of the orbits $\Gamma\theta_i$ for $i = 1, \dots, n$.*

Our main result is as follows.

Theorem 4.3. *Let P be a symmetric alcoved polytope of irreducible and reduced type $t \neq F_4$. Then P can be generated by h vertices, where h is the Coxeter number of t .*

Proof. If $\dim V = 2$, our claim follows from Lemma 3.5. The simply laced case is proved below in Theorem 4.4. The case B_n and C_n will be shown in Theorem 4.5. \square


Symmetric alcoved polytopes of type F_4 may require up to 24 generators while the Coxeter number is 12; see Theorem 4.6 for a detailed analysis.

Note that by Remark 3.4 the bound from Theorem 4.3 is the candidate for the best possible result for general (not necessarily symmetric) alcoved polytopes, since $h = |\Phi|/\dim(V)$. This works in the A_n case, as explained above. However, in the case of the root system D_4 there are exceptions of non-symmetric alcoved polytopes where more generators are needed; see Example 3.7. It is an interesting question for which types of root system the bound from Theorem 4.3 can be generalized to the non-symmetric case.

4.1. Simply laced case: types A , D , and E . As indicated before, we will prove Theorem 4.3 in several steps. We begin with the simply laced case. Recall that a root system Φ is called simply laced if all roots have the same length, i.e. if the number $\langle a, a \rangle$ is constant for all $a \in \Phi$. The types of the simply laced root systems are precisely the types A , D and E .

Theorem 4.4. *Let P be a symmetric alcoved polytope of type t , where t is irreducible, reduced and simply laced, i.e. either A_n for $n \geq 1$, D_n for $n \geq 4$ or E_6 , E_7 or E_8 . Then P can be generated by h vertices.*

Proof. We assume first that t is not E_8 . We choose a basis b_1, \dots, b_n of Φ and look at the corresponding Dynkin diagram. We number the basis elements in the following way:



Note that the numbering in the D_n case is opposite to the numbering often found in the literature. Recall that every vertex in the Dynkin diagram represents a basis element b_i and that $\langle b_i, b_j \rangle$ is $-\langle b_j, b_i \rangle$ if the vertices for b_i and b_j are connected, and 0, if they are not. Since the Dynkin diagram is a tree, for all i and j there exists a geodesic $[b_i, b_j]$ connecting the vertices b_i and b_j . We first show the following claim.

Claim: Let $i \geq 1$, if t is A_n , and $i \geq 2$, if t is D_n , and $i \geq 3$, if t is E_6 or E_7 . Then for all $j \geq i$ we have

$$s_j \circ \dots \circ s_{i+1}(b_i) = \sum_{\nu=i}^j b_\nu.$$

Let us prove the claim by induction on j . If j is equal to i there is nothing to prove. Assume that the claim holds for some $j \geq i$ with $j < n$. Then we calculate

$$\begin{aligned}
s_{j+1} \circ s_j \circ \dots \circ s_{i+1}(b_i) &= s_{j+1}\left(\sum_{\nu=i}^j b_\nu\right) \\
&= \sum_{\nu=i}^{j-1} b_\nu + s_{j+1}(b_j) \\
&= \sum_{\nu=i}^{j-1} b_\nu + b_j + b_{j+1} \\
&= \sum_{\nu=i}^{j+1} b_\nu,
\end{aligned}$$

which finishes the proof of the claim.

Recall that for all $i = 1, \dots, n$ we have

$$\theta_i = s_n \circ \dots \circ s_{i+1}(b_i).$$

From the claim we deduce that $\theta_i = \sum_{\nu=i}^n b_\nu$, which implies

$$(*) \quad \theta_i = \sum_{b_\nu \in [b_i, b_n]} b_\nu$$

except possibly in the cases $t = D_n$ and $i = 1$ or $t = E_6$ or E_7 and $i = 1, 2$. It can easily be checked directly that the statement $(*)$ also holds in these remaining cases.

Since in all cases b_n is a neighbour of b_{n-1} in the Dynkin diagram, $(*)$ implies that

$$\theta_i = b_n + \sum_{b_\nu \in [b_i, b_{n-1}]} b_\nu.$$

In a simply laced root system, the Weyl group acts transitively on the set of roots, see [Bou, VI, 1.3, Proposition 11]. Hence every symmetric alcoved polytope is of the form

$$P_\lambda = \bigcap_{a \in \Phi} \{x : \langle a, x \rangle \leq \lambda\}$$

for some $\lambda > 0$.

Since b_1, \dots, b_n is a basis of V , there is a unique point $x \in V$ such that

$$\langle b_n, x \rangle = \lambda, \langle b_{n-1}, x \rangle = \dots = \langle b_2, x \rangle = \langle b_1, x \rangle = 0.$$

We will show now that x is a vertex of P_λ .

Note that the highest root in Φ is of the following form:

$$\begin{aligned}
A_n : & \quad b_1 + \dots + b_n \\
D_n : & \quad b_1 + b_2 + 2b_3 + \dots + 2b_{n-1} + b_n \\
E_6 : & \quad b_1 + 2b_2 + 3b_3 + 3b_4 + 2b_5 + b_6 \\
E_7 : & \quad 2b_1 + 2b_2 + 3b_3 + 4b_4 + 3b_5 + 2b_6 + b_7
\end{aligned}$$


Therefore for every positive root $a \in \Phi$ we find that $\langle a, x \rangle = 0$ or $\langle a, x \rangle = \langle b_n, x \rangle = \lambda$. For the negative roots there is nothing to do, since they assume a non-positive value on x . Hence we find that x lies in P_λ . Since $\theta_1, \dots, \theta_n$ are independant (see [Bou], VI, 1.11, Proposition 33), and x lies in all hyperplanes $\{\langle \theta_i, - \rangle = \lambda\}$, we find that x is a vertex.

Since the Weyl group maps faces to faces, the elements $\omega^k x$ for $k = 0, \dots, h-1$ are also vertices of P_λ , where $\omega = s_{b_1} \circ \dots \circ s_{b_n}$ is the corresponding Coxeter element. Since x satisfies $\langle \theta_i, x \rangle = \lambda$ for all $i = 1, \dots, n$, the vertex $y = \omega^k(x)$ satisfies $\langle \omega^k \theta_i, y \rangle = \lambda$ for all $i = 1, \dots, n$. Since every root is of the form $\omega^k \theta_i$ for suitable i and k , every hyperplane appearing at the boundary of P_λ contains one of the vertices $\omega^k x$ for $k = 0, \dots, h-1$. This implies our claim for all types except possibly E_8 .

We conclude the proof by discussing the type E_8 . Let

$$B = \begin{pmatrix} 1/2 & 1 & -1 & 0 & 0 & 0 & 0 & 0 \\ -1/2 & 1 & 1 & -1 & 0 & 0 & 0 & 0 \\ -1/2 & 0 & 0 & 1 & -1 & 0 & 0 & 0 \\ -1/2 & 0 & 0 & 0 & 1 & -1 & 0 & 0 \\ -1/2 & 0 & 0 & 0 & 0 & 1 & -1 & 0 \\ -1/2 & 0 & 0 & 0 & 0 & 0 & 1 & -1 \\ -1/2 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1/2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad \omega = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & -1 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & -1 \end{pmatrix}, \quad \Theta = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{pmatrix}.$$

Then the columns b_1, b_2, \dots, b_8 of the matrix B are a basis for a root system of type E_8 . It has the following Dynkin diagram




The matrix ω represents the Coxeter element associated to b_1, \dots, b_8 , and the columns of Θ are the roots $\theta_i = s_n \circ \dots \circ s_{i+1}(b_i)$ expressed in the basis b_1, \dots, b_8 . Let $x = \frac{1}{2}(b_2 + b_3)$ in V . We verify by explicit calculation that the point x lies on the root hyperplanes corresponding to $\omega^{17}\theta_1, \omega^{20}\theta_1, \omega^{16}\theta_2, \omega^{18}\theta_2, \omega^{17}\theta_3, \omega^{15}\theta_4, \omega^{20}\theta_4, \omega^9\theta_5, \omega^{14}\theta_5, \omega^{13}\theta_6, \omega^{12}\theta_7, \omega^{11}\theta_8, \omega^{20}\theta_8$, and $\omega^{25}\theta_8$. Hence x touches all orbits of the root hyperplanes under the Coxeter element ω , so the orbit of x forms a 30-element generating set for the symmetric alcoved polytope P_1 of E_8 . Modulo scaling, there is only one symmetric alcoved polytope of type E_8 , which has f -vector (19440, 207360, 483840, 483840, 241920, 60480, 6720, 240, 1), computed using Polymake [Ga-Jo]. \square

4.2. Types B and C . Since the symmetric alcoved polytopes of type C_n coincide with the symmetric alcoved polytopes of type B_n , it suffices to consider the B_n case. Here we may assume $V = \mathbb{R}^n$ and $\Phi = \{\pm a_i : i = 1, \dots, n\} \cup \{\pm a_i \pm a_j : 1 \leq i < j \leq n\}$, where a_i is the canonical basis of V . The Coxeter number for types B_n and C_n is $2n$.

Theorem 4.5. *Let P be a symmetric alcoved polytope of type B_n or C_n for $n \geq 2$. Then P can be generated by $2n$ vertices.*

Proof. We choose as a basis $b_1 = a_1 - a_2, b_2 = a_2 - a_3, \dots, b_{n-1} = a_{n-1} - a_n$ and $b_n = a_n$ of Φ . The Dynkin diagram of type B_n is the following:



We then calculate the roots $\theta_1, \dots, \theta_n$ as

$$\theta_n = a_n \text{ and } \theta_i = a_n + a_i \text{ for } i = 1, \dots, n-1.$$

Recall that the union of the orbits of the θ_i under the group Γ generated by the Coxeter element $\omega = s_1 \circ \dots \circ s_n$ is the whole root system. Since Weyl group elements map long roots to long roots and short roots to short roots, we deduce that the set of short roots is equal to Γb_n , and that the set of long roots is equal to $\Gamma b_1 \cup \dots \cup \Gamma b_{n-1}$.

A type B_n or C_n symmetric alcoved polytope P is of the form

$$P_{\lambda, \mu} = \bigcap_i \{x : \langle \pm a_i, x \rangle \leq \mu\} \cap \bigcap_{i < j} \{x : \langle (\pm a_i \pm a_j), x \rangle \leq \lambda\}$$

for positive numbers λ and μ satisfying $\mu \leq \lambda \leq 2\mu$. In the extreme cases $\mu = \lambda$ and $\lambda = 2\mu$, only long or short root hyperplanes appear as facets, and the others support lower dimensional faces. However, this would not affect our reasoning. Let x be the point in V defined by

$$\langle a_1, x \rangle = \dots = \langle a_{n-1}, x \rangle = \lambda - \mu \text{ and } \langle a_n, x \rangle = \mu.$$

Then x is a point in $P_{\lambda, \mu}$ lying in the n independent hyperplanes $\{\langle \theta_n, - \rangle = \mu\}$ and $\{\langle \theta_i, - \rangle = \lambda\}$ for $i = 1, \dots, n-1$. Therefore x is a vertex. As in the proof of Theorem 4.4, the h vertices $\omega^k x$ for $k = 0, \dots, h-1$ touch every boundary hyperplane. Hence they form a generating set of cardinality h . \square

4.3. Type F_4 . Let $V = \mathbb{R}^4$ with canonical basis a_1, \dots, a_4 . Then

$$\Phi = \{\pm a_i : i = 1, \dots, 4\} \cup \{\pm a_i \pm a_j : 1 \leq i < j \leq 4\} \cup \left\{ \frac{1}{2}(\pm a_1 \pm a_2 \pm a_3 \pm a_4) \right\}$$

is a root system of type F_4 in V . It consists of 24 long roots (of the form $\pm a_i \pm a_j$) and 24 short roots.

A symmetric alcoved polytope of type F_4 is of the form

$$P_{\lambda, \mu} = \bigcap_i \{ \langle \pm a_i, x \rangle \leq \mu \} \cap \{ \langle (\pm a_1 \pm a_2 \pm a_3 \pm a_4), x \rangle \leq 2\mu \} \cap \bigcap_{i < j} \{ \langle (\pm a_i \pm a_j), x \rangle \leq \lambda \}$$

for positive numbers λ and μ satisfying $\mu \leq \lambda \leq 2\mu$. This condition ensures that all root hyperplanes are support hyperplanes. When $\mu = \lambda$ or $\lambda = 2\mu$, either only long roots or only short roots appear as facet normals and the others support faces of lower dimension. When $\mu < \lambda < 2\mu$, all root hyperplanes support facets.

The convex hull of short roots of F_4 is a regular polytope called the *24-cell*. The long roots are the vertices of its dual. The 24-cell is combinatorially equivalent to $P_{1,1}$. The root system of type F_4 is self-dual. By doubling the lengths of the short roots, and then rescaling everything by $\frac{1}{\sqrt{2}}$, we get back F_4 , rotated, with the roles of long and short roots

switched. More concretely, the matrix $A = \frac{1}{2} \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & -1 \end{pmatrix}$ maps the long roots to short roots and the doubled short roots to long roots. Then A^{-T} is a linear isomorphism from

$P_{\lambda,\mu}$ to $P_{2\mu,\lambda}$, so these two polytopes have exactly the same combinatorics, in particular, the same number of generators.


Theorem 4.6. *Let $P_{\lambda,\mu}$ be an alcoved polytope of type F_4 , using the notation above.*

(1) *If $\frac{4}{3}\mu \leq \lambda \leq \frac{3}{2}\mu$, then $P_{\lambda,\mu}$ is generated by 12 elements.*

(2) *If $\mu \leq \lambda \leq \frac{4}{3}\mu$ or $\frac{3}{2}\mu \leq \lambda \leq 2\mu$, then $P_{\lambda,\mu}$ is generated by 24 elements.*

In each of these cases, if the inequalities among the λ and μ are strict, then the numbers of generators, 12 and 24 respectively, are minimum possible.

Proof. The Dynkin diagram of a root system of type F_4 is the following:



with the basis $b_1 = a_2 - a_3, b_2 = a_3 - a_4, b_3 = a_4$ and $b_4 = \frac{1}{2}(a_1 - a_2 - a_3 - a_4)$ of Φ . We calculate the roots $\theta_1, \dots, \theta_4$ associated to the basis $\{b_1, b_2, b_3, b_4\}$:

$$\begin{aligned} \theta_1 &= a_1 - a_3 \\ \theta_2 &= a_1 - a_2 \\ \theta_3 &= \frac{1}{2}(a_1 - a_2 - a_3 + a_4) \\ \theta_4 &= \frac{1}{2}(a_1 - a_2 - a_3 - a_4) \end{aligned}$$

Case (1): The alcoved polytope is of the form $P_{\lambda,\mu}$ with $\frac{4}{3}\mu \leq \lambda \leq \frac{3}{2}\mu$. Let x be the element of V given by

$$\langle a_1, x \rangle = 2\lambda - 2\mu, \langle a_2, x \rangle = \lambda - 2\mu, \langle a_3, x \rangle = \lambda - 2\mu, \langle a_4, x \rangle = 0.$$

We check that x is a point in $P_{\lambda,\mu}$ as follows. Using $\mu < \lambda$ and $2\lambda \leq 3\mu$, we find

$$-\mu \leq 2\lambda - 2\mu \leq \mu \text{ and } -\mu \leq \lambda - 2\mu \leq \mu.$$

Hence x lies in $\bigcap_i \{\langle \pm a_i, x \rangle \leq \mu\}$. Moreover, we have

$$\pm(2\lambda - 2\mu) \pm (\lambda - 2\mu) \pm (\lambda - 2\mu) \leq 2\lambda - 2\mu + 2(2\mu - \lambda) = 2\mu,$$

so x lies in $\{\langle (\pm a_1 \pm a_2 \pm a_3 \pm a_4), x \rangle \leq 2\mu\}$. Checking the long roots we find

$$\pm(2\lambda - 2\mu) \pm (\lambda - 2\mu) \leq 2\lambda - 2\mu + 2\mu - \lambda = \lambda, \text{ and}$$

$$-\lambda \leq 2\lambda - 2\mu \leq \lambda \text{ and } -\lambda \leq \lambda - 2\mu \leq \lambda.$$

It remains to check

$$\pm(\lambda - 2\mu) \pm (\lambda - 2\mu) \leq 4\mu - 2\lambda \leq \lambda,$$

which follows from the condition $4\mu \leq 3\lambda$.

Then we calculate $\langle \theta_1, x \rangle = \lambda, \langle \theta_2, x \rangle = \lambda, \langle \theta_3, x \rangle = \mu, \langle \theta_4, x \rangle = \mu$. Hence x lies in four linearly independent facets of $P_{\lambda,\mu}$, so it is a vertex. By the same argument as in the proof of Theorem 4.4, the vertices $\omega^k x$ for $k = 0, \dots, h-1$ generate $P_{\lambda,\mu}$.

A computation using Gfan [Jen] shows that all values of λ and μ satisfying $\frac{4}{3}\mu < \lambda < \frac{3}{2}\mu$ give the same combinatorial type for $P_{\lambda,\mu}$. A Polymake [Ga-Jo] computation shows that the polytope is simple in this case, i.e. each vertex lies in exactly 4 facets. Hence at least 12 vertices are necessary to meet all 48 facets. The polytope has f -vector $(288, 576, 336, 48, 1)$.

Case (2): The remaining cases are $\mu \leq \lambda < \frac{4}{3}\mu$ and $\frac{3}{2}\mu < \lambda \leq 2\mu$. However, these two cases are equivalent via the isomorphism of $P_{\lambda,\mu}$ and $P_{2\mu,\lambda}$ seen above.

Let us consider the case $\frac{3}{2}\mu < \lambda \leq 2\mu$. The point $x_1 = (\mu, \mu - \lambda, \lambda - 2\mu, 0)$ lies on the facets perpendicular to θ_2, θ_3 , and θ_4 , while the point $x_2 = (\mu, \lambda - 2\mu, \mu - \lambda, 0)$ lies on the facets perpendicular to θ_1, θ_3 , and θ_4 . They also lie in the hyperplane defined by $\langle a_1, x \rangle = \mu$, so they are vertices of the polytope. Since $x_1, x_2 \in P_{\lambda,\mu}$ and together they touch all hyperplanes perpendicular to θ_i 's, the union of the orbits of x_1 and x_2 under the group Γ forms a generating set of size at most 24.

Computation by Gfan [Jen] shows that all values of λ and μ satisfying $\frac{3}{2}\mu < \lambda < 2\mu$ give the same the combinatorial type for $P_{\lambda,\mu}$, which is simple and has f -vector $(192, 384, 240, 48, 1)$. The 192 vertices of $P_{\lambda,\mu}$ are obtained from x_1 (or equivalently x_2) by permuting coordinates in any of the 24 ways and changing signs in any of the 8 ways, for a total of 192 distinct combinations. It can then be seen directly that each vertex lies in exactly one long-root facet and there are 24 long-root facets, so $P_{\lambda,\mu}$ needs at least 24 generators. \square

REFERENCES

- [Bou] N. Bourbaki: *Lie groups and Lie algebras*, Chapter 4-6. Translated from the 1968 French original by Andrew Pressley. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2002.
- [De-St] M. Develin, B. Sturmfels: *Tropical convexity*. Doc. Math. 9 (2004) 1-27.
- [Ga-Jo] E. Gawrilow and M. Joswig: *polymake: a framework for analyzing convex polytopes*. Polytopes – combinatorics and computation (Oberwolfach, 1997), 4373.
- [Jen] A. N. Jensen: *Gfan, a software system for Gröbner fans and tropical varieties*. Available at <http://home.imf.au.dk/jensen/software/gfan/gfan.html>
- [Jo-Ku] M. Joswig, K. Kulas: *Tropical and Ordinary Convexity combined*. Adv. Geometry 10 (2010) 333-352.
- [La-Po] T. Lam, A. Postnikov: *Alcoved Polytopes I*. Discrete Comput. Geom. 38 (2007) 453-478.
- [Pa] S. Payne: *Lattice polytopes cut out by root systems and the Koszul property*. Adv. in Math. 220 (2009) 926-935.

INSTITUT FÜR MATHEMATIK, GOETHE-UNIVERSITÄT FRANKFURT, ROBERT-MAIER-STR., 6-8, D-60325 FRANKFURT-AM-MAIN, GERMANY

E-mail address: `werner@math.uni-frankfurt.de`

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, 686 CHERRY ST, ATLANTA GA 30332, USA

E-mail address: `jyu@math.gatech.edu`